



Contents lists available at ScienceDirect

Journal of Cardiology Cases

journal homepage: www.elsevier.com/locate/jccase

Case Report

A rare case of cardiac metastasis from uterine cervical adenocarcinoma

Shokoufeh Hajsadeghi, MD^a, Marziyeh Pakbaz, MD^{b,*}, Alireza Sadeghipour, MD^c,
Sam Zeraatian nejad davani^d, Ali Shamsedini, MD^e

^a Professor of Cardiology, Research Center for Prevention of Cardiovascular Disease, Institute of Endocrinology & Metabolism, Iran University of Medical Sciences, Tehran, Iran

^b Assistant Professor of Cardiology, Department of Cardiovascular Disease, Hazrat-e Rasool General Hospital, Iran University of Medical Sciences, Sattar-khan Avenue, Niyayesh Street, 1445613131 Tehran, Iran

^c Professor of Pathology, Department of Pathology and Oncopathology Research Center, Iran University of Medical Sciences, Tehran, Iran

^d Assistant Professor of Cardiac Surgery, Department of Cardiac Surgery, Hazrat-e Rasool General Hospital, Iran University of Medical Sciences, Tehran, Iran

^e Resident of Cardiology, Department of Cardiovascular Disease, Hazrat-e Rasool General Hospital, Iran University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Article history:

Received 28 September 2020

Revised 13 November 2020

Accepted 24 November 2020

Available online xxx

Keywords:

Uterine cervical neoplasms

Adenocarcinoma

Cardiac metastasis

Cardiac mass

Uterine cervix

ABSTRACT

Metastatic cardiac tumors are rare and are usually from lung, breast, and esophageal cancers. Although uterine cervical cancer is relatively common, its metastasis to the heart is extremely rare. Herein, we report an unusual case of metastatic cervical adenocarcinoma to the heart presenting with a huge right atrial mass and tamponade. The cardiac mass was surgically resected and pathologic study with immunohistochemistry staining confirmed the diagnosis.

Learning objective: Metastatic cardiac tumors are rare. They usually originate from lung, breast, and esophageal cancers and mostly involve the pericardium and myocardium. The endocardium is rarely affected. Cervical cancer is one of the most common gynecologic cancers with a relatively favorable prognosis unless it metastasizes to distant organs. Metastases to the heart are rarely reported. The patient may be asymptomatic or present with hemodynamic instability due to tamponade state and/or a huge intracardiac mass.

© 2020 Japanese College of Cardiology. Published by Elsevier Ltd. All rights reserved.

Introduction

Metastatic cardiac tumors are rare with a reported incidence of 1.2% based on autopsy studies [1]. The incidence may be underestimated, however, because these lesions are often clinically silent and found incidentally during clinical imaging for other purposes or may be unrecognized until an autopsy study is performed [2]. Theoretically, any malignant tumor is capable to metastasize to the heart; the most common ones include lung, breast, and esophageal cancers. Metastases from infradiaphragmatic organs are less frequent [3]. Herein, we report an unusual case of metastatic cervical cancer to the heart presenting with a huge right atrial mass and tamponade.

Case report

The patient was a 60-year-old female, a known case of endometrial cancer of the uterine cervix who presented at our center with progressive dyspnea. Six months previously, after being

evaluated for abnormal vaginal bleeding, she had been found to have a stage IIb-infiltrative cervical mass with extension to the proximal vagina and also to the adjacent parametrium revealed by abdominopelvic sonography and magnetic resonance imaging (MRI). Cervical biopsy with immunohistochemistry (IHC) study reported at another center had been strongly positive for P53 and P16 markers, and somewhat positive for CK7, CK20, and CDX2, compatible with the usual type of primary endocervical adenocarcinoma. She had undergone chemoradiation for several months but we were not informed about the details of the treatment regimen. A control MRI some weeks before presentation, had shown a slight shrinkage of the mass. Total hysterectomy had been planned but at the same time, she developed progressive dyspnea. By arrival at our center, the blood pressure was 95/60 mmHg and the heart rate was 125/min. On physical examination there were muffled heart sounds and jugular vein distension. A pulsus paradoxus of 25 mmHg was also noted. She experienced a rapid deterioration of clinical symptoms including massive ascites and at rest dyspnea. The patient was clinically at tamponade state. Transthoracic and transesophageal echocardiography revealed a huge mass almost completely occupying the right atrium with attachment to the intra-atrial septum and tricuspid valve (TV) leaflets (Fig. 1).

* Corresponding author.

E-mail address: marzi.pakbaz@gmail.com (M. Pakbaz).

